

# The MEG experiment at PSI: search for the $\mu^+ \rightarrow e^+ \gamma$ decay

---

Giovanni Signorelli

INFN Pisa and Pisa University (Italy)  
for the MEG collaboration



NEW TRENDS IN HIGH ENERGY PHYSICS

Yalta, Ukraine - September 16-23, 2006





**WASEDA**

**UCIrvine**



PAUL SCHERRER INSTITUT  
**PSI**

### **Univ. of Tokyo**

Y. Hisamatsu, T. Iwamoto, S. Mihara, **T. Mori**, H. Natori, H. Nishiguchi, W. Ootani, R. Sawada, S. Yamashita, Y. Uchiyama

### **KEK**

T. Haruyama, K. Kasami, A. Maki, Y. Makida, A. Yamamoto

### **Waseda Univ.**

T. Doke, J. Kikuchi, S. Suzuki, K. Terasawa

### **INFN and University of Pisa**

**A. Baldini**, C. Bemporad, F. Cei, M. Grassi, D. Nicolò, A. Papa, R. Pazzi, F. Sergiampietri, G. Signorelli

### **INFN and Univ. of Genova**

S. Dussoni, F. Gatti, R. Valle

### **INFN and Univ. of Pavia**

G. Boca, P. W. Cattaneo, G. Cecchet, A. De Bari, R. Nardò, M. Rossella

### **INFN and Univ. of Roma I**

A. Barchiesi, D. Zanello

### **INFN and Univ. of Lecce**

M. Panareo

### **Paul Scherrer Institute**

J. Egger, M. Hildebrandt, P.-R. Kettle, **S. Ritt**, M. Schneebeli

### **BINP Novosibirsk**

L. M. Barkov, A. A. Grebenuk, D. N. Grigoriev, B. I. Khazin, N. M. Ryskulov

### **JINR Dubna**

A. Korenchenko, N. Kravchuk, A. Moiseenko, D. Mzavia

### **Univ. of California, Irvine**

W. Molzon, M. Hebert, P. Huwe, J. Perry, V. Tumakov, F. Xiao, S. Yamada

# Outline

- Physics **motivation** for a  $\mu \rightarrow e\gamma$  experiment
- The  $\mu \rightarrow e\gamma$  decay
- The **detector**
  - Beam line & target
  - Spectrometer
  - Timing Counter
  - LXe calorimeter
  - Calibrations
  - Electronics
- **Status**
- Future

# The SM. And beyond...


- The Standard Model is made of particles + interactions + symmetries
  - Physics symmetries (Lorentz, Poincare'...)
  - Gauge symmetries (charge, chirality)
  - Accidental symmetries

$B$	$L$	$L_i$
baryon number	lepton number	lepton family number
$p \rightarrow e^- \pi^0$	$\nu$ masses	$\mu \rightarrow e\gamma$

- These symmetries are called accidental because there is no general rule that imposes them - they just “happen” to be in the SM
- The research for the failure of one of these symmetries could shed new light on particle physics
- Three complementary searches to probe the Standard Model.

# The $\mu \rightarrow e\gamma$ decay

- The  $\mu \rightarrow e\gamma$  decay is **forbidden** in the **SM** because of the (accidental) conservation of **lepton family numbers**
- The introduction of **neutrino masses and mixings** induces  $\mu \rightarrow e\gamma$  radiatively, but at a **negligible level**




$$\Gamma(\mu \rightarrow e\gamma) \approx \underbrace{\frac{G_F^2 m_\mu^5}{192\pi^3}}_{\mu - \text{decay}} \underbrace{\left(\frac{\alpha}{2\pi}\right)}_{\gamma - \text{vertex}} \underbrace{\sin^2 2\theta \sin^2 \left(\frac{1.27\Delta m^2}{M_W^2}\right)}_{\nu - \text{oscillation}}$$

$$\approx 10^{-53}$$


- All **SM extensions enhance the rate** through **mixing** in the **high energy sector** of the theory

# For instance... predictions





- **SUSY SU(5)** predictions: LFV induced by finite slepton mixing through radiative corrections. The mixing could be large due to the top-quark mass at a level of  $10^{-12} - 10^{-15}$
- **SO(10)** predicts even larger BR:
  - $m(\tau)/m(\mu)$  enhancement
- Models with **right-handed neutrinos** also predict large BR
- $\Rightarrow$  clear evidence for physics beyond the SM.

R. Barbieri et al., Nucl. Phys. B445 (1995) 215  
 J. Hisano et al., Phys. Lett. B391 (1997) 341  
 R. Ciafaloni, A. Romanino, A. Strumia, Nucl. Phys. B458 (1996)  
 J. Hisano, N. Nomura, Phys. Rev. D59 (1999)




# Historical perspective



Each improvement linked to an  
**improvement in the technology**

$$N_{\text{sig}} = R_\mu T \frac{\Delta\Omega}{4\pi} \epsilon_\gamma \epsilon_e \epsilon_{\text{cut}} B_{\mu \rightarrow e\gamma}$$



# Signal and Background



Exp./Lab	Year	$\Delta E_e/E_e$ (%)	$\Delta E_\gamma/E_\gamma$ (%)	$\Delta t e\gamma$ (ns)	$\Delta \theta e\gamma$ (mrad)	Stop rate ( $s^{-1}$ )	Duty cyc. (%)	BR (90% CL)
SIN	1977	8.7	9.3	1.4	-	$5 \times 10^5$	100	$3.6 \times 10^{-9}$
TRIUMF	1977	10	8.7	6.7	-	$2 \times 10^5$	100	$1 \times 10^{-9}$
LANL	1979	8.8	8	1.9	37	$2.4 \times 10^5$	6.4	$1.7 \times 10^{-10}$
Crystal Box	1986	8	8	1.3	87	$4 \times 10^5$	(6..9)	$4.9 \times 10^{-11}$
MEGA	1999	1.2	4.5	1.6	17	$2.5 \times 10^8$	(6..7)	$1.2 \times 10^{-11}$
MEG	2006	0.8	4	0.15	19	$2.5 \times 10^7$	100	$1 \times 10^{-13}$

# MEG experimental method

Easy signal selection with  $\mu^+$  at rest




- Stopped beam of  $>10^7 \mu$  /sec in a  $150 \mu\text{m}$  target
- $\gamma$  detection  
Liquid Xenon calorimeter based on the scintillation light
  - fast: 4 / 22 / 45 ns
  - high LY:  $\sim 0.8 * \text{NaI}$
  - short  $X_o$ : 2.77 cm
- $e^+$  detection  
magnetic spectrometer composed by solenoidal magnet and drift chambers for momentum scintillation counters for timing

# Beam line & target

$\pi E_5$  beam line at PSI



Optimization of the beam elements:

- Muon momentum ~ **29 MeV/c**
- Wien filter for  **$\mu/e$  separation**
- Solenoid to couple beam and spectrometer (**BTS**)
- **Degrader** to reduce the momentum for a **150  $\mu\text{m}$**  target




Present results (1.8 mA):



- $R_\mu$  (total)  $1.3 \times 10^8 \mu^+/\text{s}$
- $R_\mu$  (after Triplet 2)  $9.4 \times 10^7 \mu^+/\text{s}$
- $\mu/e$  separation  $11.8 \text{ cm} (7.2 \sigma)$
- $R_\mu$  (exp. on target)  $6.4 \times 10^7 \mu^+/\text{s}$
- $\mu$  spot (exp. on target)  $\sigma_V \approx \sigma_H \approx 10 \text{ mm}$



# COBRA spectrometer





Non uniform  
magnetic field  
decreasing from  
the center to  
the periphery





Compensation  
coil for LXe  
calorimeter

$$|\vec{B}| < 50 \text{ G}$$




# Positron Tracker



- 
- 16 chambers radially aligned with 10° intervals
  - 2 staggered arrays of drift cells
  - 1 signal wire and 2 x 2 vernier cathode strips made of 15 µm kapton foils and 0.45 µm aluminum strips
  - Chamber gas: He-C<sub>2</sub>H<sub>6</sub> mixture

transverse coordinate (t drift)





longitudinal coordinate (charge division)

Measurements at Tokyo University:

$$\sigma_R = 93 \pm 10 \mu\text{m}$$

$$\sigma_Z = 425 \pm 7 \mu\text{m}$$

# Drift chambers




- Full scale test in November
- Summary of Drift Chamber simulation

$$\delta P_{e^+} / P_{e^+} = 0.7 \div 0.9\%$$


$$\delta\theta_{e^+} = 9 \div 12 \text{ mrad}$$

$$\delta x_{orig} = 2.1 \div 2.5 \text{ mm}$$



FWHM



# Timing Counter




- Two layers of scintillators:  
Outer layer, read out by PMTs: timing measurement  
Inner layer, read out with APDs at  $90^\circ$ : z-trigger
- Obtained goal  $\sigma_{\text{time}} \sim 40 \text{ psec}$  (100 ps FWHM)




Exp. application (*)	Counter size (cm) (T x W x L)	Scintillator	PMT	$\lambda_{\text{att}}$ (cm)	$\sigma_t(\text{meas})$	$\sigma_t(\text{exp})$
G.D.Agostini	3 x 15 x 100	NE114	XP2020	200	120	60
T. Tanimori	3 x 20 x 150	SCSN38	R1332	180	140	110
T. Sugitate	4 x 3.5 x 100	SCSN23	R1828	200	50	53
R.T. Gile	5 x 10 x 280	BC408	XP2020	270	110	137
TOPAZ	4.2 x 13 x 400	BC412	R1828	300	210	240
R. Stroynowski	2 x 3 x 300	SCSN38	XP2020	180	180	420
Belle	4 x 6 x 255	BC408	R6680	250	90	143
<b>MEG</b>	<b>4 x 4 x 90</b>	<b>BC404</b>	<b>R5924</b>	<b>270</b>	<b>38</b>	

Best existing TC

# Timing Counter



- Two layers of scintillators:  
Outer layer, read out by PMTs: timing measurement  
Inner layer, read out with APDs at  $90^\circ$ : z-trigger
- Obtained goal  $\sigma_{\text{time}} \sim 40$  psec (100 ps FWHM)





Exp. application (*)	Counter size (cm) (T x W x L)	Scintillator	PMT	$\lambda_{\text{att}}$ (cm)	$\sigma_t(\text{meas})$	$\sigma_t(\text{exp})$
G.D.Agostini	3 x 15 x 100	NE114	XP2020	200	120	60
T. Tanimori	3 x 20 x 150	SCSN38	R1332	180	140	110
T. Sugitate	4 x 3.5 x 100	SCSN23	R1828	200	50	53
R.T. Gile	5 x 10 x 280	BC408	XP2020	270	110	137
TOPAZ	4.2 x 13 x 400	BC412	R1828	300	210	240
R. Stroynowski	2 x 3 x 300	SCSN38	XP2020	180	180	420
Belle	4 x 6 x 255	BC408	R6680	250	90	143
MEG	4 x 4 x 90	BC404	R5924	270	38	

Best existing TC


# The calorimeter

- $\gamma$  Energy, position, timing
- Homogeneous  $0.8 \text{ m}^3$  volume of liquid Xe
  - 10 % solid angle
  - $65 < r < 112 \text{ cm}$
  - $|\cos\theta| < 0.35 \quad |\phi| < 60^\circ$
- Only scintillation light
- Read by 848 PMT
  - 2" photo-multiplier tubes
  - Maximum coverage FF (6.2 cm cell)
  - Immersed in liquid Xe
  - Low temperature (165 K)
  - Quartz window (178 nm)
- Thin entrance wall
- Singularly applied HV
- Waveform digitizing @ 2 GHz
  - Pileup rejection




# Xe properties

- Fast
  - $\tau_{\text{singlet}} = 4.2 \text{ ns}$
  - $\tau_{\text{triplet}} = 22 \text{ ns}$
  - $\tau_{\text{recomb}} = 45 \text{ ns}$
- Particle ID
  - LY alpha =  $1.2 \times \text{LY gamma/e}$
- High LY ( $\approx \text{NaI}$ )
  - 40000 phe/MeV
- $n = 1.65$
- $Z=54, \rho=2.95 \text{ g/cm}^3 (X_O=2.7 \text{ cm}), R_M=4.1 \text{ cm}$
- No self-absorption ( $\lambda_{\text{Abs}}=\infty$ )





...in progress




# The LXe calorimeter prototype

- A 100 liters, 240 PMTs **large prototype** was built and extensively tested to demonstrate the calorimeter performance
- **$\alpha$ -sources** and **LEDs** used for PMT calibrations and monitoring





# LXe calorimeter R&D



- Energy **resolution** strongly depends on **absorption**. A long R&D to insure  $L(\text{Abs}) > 3$  m with a circulation/**purification** system
- Measurement of **energy and timing resolution** with high energy photons: 55 MeV photons from pion charge exchange reaction




Lab Frame




Two tests in **2003** and **2004** demonstrated the calibration procedure and the resolutions








- In the **back-to-back** raw spectrum we see the **correlation**
  - $83 \text{ MeV} \Leftrightarrow 55 \text{ MeV}$
  - The  $129 \text{ MeV}$  line is visible in the NaI because Xe is sensitive to neutrons ( $9 \text{ MeV}$ )



# Resolutions @ 55 MeV





- Select negative pions in the beam
- $65 \text{ MeV} < E(\text{NaI}) < 95 \text{ MeV}$
- Collimator cut ( $r < 4 \text{ cm}$ )



# Position dependence


- small FWHM residual dependence
- no significant peak shift
- The resolution is always better than 5% FWHM

4.8%	4.6%
4.5%	5.0%







# MEG (LXe) calibrations

- A reliable result depend on a constant **calibration** and **monitoring** of the apparatus
  - alpha Sources (on wires and wall)
  - Proton accelerator  ${}^7\text{Li}(p, \gamma_{17.6}){}^8\text{Be}$  design under way
  - Neutron generator  ${}^{58}\text{Ni}(n, \gamma_9){}^{59}\text{Ni}$
  - Charge exchange reaction (Panofsky)  
$$\begin{aligned} \pi^- p &\rightarrow \pi^0 n \\ \pi^0 &\rightarrow \gamma\gamma \end{aligned}$$
- Calibration frequency is different




# Alpha sources


- 4 W wires suspended inside the liquid xenon
  - unique to liquid
  - Po and Am sources
- QEs determined by **comparison** of alpha source signal in cold gaseous xenon and **MC** determined at a 10% level



27



First Ver.



Second Ver.






Final Ver.

# Proton accelerator

- Positioned opposite to the muon beam
- Essentially two reactions



<i>Reaction</i>	<i>Resonance energy</i>	$\sigma$ peak	$\gamma$ -lines
$Li(p,\gamma)Be$	440 keV	5 mb	17.6 MeV, 14.6 MeV
$B(p,\gamma)C$	163 keV	$2 \cdot 10^{-1}$ mb	4.4 MeV, 11.7 MeV, 16.1 MeV

- tested at Legnaro (Italy) van de Graaf with a large NaI



# Neutron/Ni $\gamma$ -line

- Reaction of thermal neutrons on Ni
- AmBe source inside a polyethylene/nickel sandwich (Cf)
- Tested with large NaI
- Will be placed behind the calorimeter




# Trigger Electronics

- 100 MHz **waveform digitizer** on VME boards that perform online pedestal subtraction
- Uses :
  - $\gamma$  energy
  - $e^+ - \gamma$  time coincidence
  - $e^+ - \gamma$  collinearity
- Built on a FADC-FPGA architecture
- More performing algorithms could be implemented




- Prototype system has been **successfully tested** on the LP
- Design of the final system is in progress
  - $\pi^0$  data
  - Charge spectrum
  - Only 32 PMT

- ❖ Beam rate  $10^8 s^{-1}$
- ❖ Fast LXe energy sum  $> 45\text{MeV}$   
 $2 \times 10^3 s^{-1}$
- gamma interaction point (PMT of max charge)
- $e^+$  hit point in timing counter
- ❖ time correlation  $\gamma - e^+$   $200 s^{-1}$
- ❖ angular correlation  $\gamma - e^+$  **20 s<sup>-1</sup>**




# Readout electronics

2.5 GHz Waveform digitization for all channels







DRS2 chip (Domino Ring Sampler)

- Custom sampling chip designed at PSI
- 2.5 GHz sampling speed @ 40 ps timing resolution
- Sampling depth 1024 bins for 8 channels/chip
- Data taken in charge exchange test to study pile-up rejection algorithms



Original



# MEG sensitivity



- Computation of the **sensitivity** based on the measured **resolutions**

FWHM $E_\gamma/E_\gamma$	5 %
FWHM $E_e/E_e$	0.9 %
$\delta t_{e\gamma}$	105 ps
$\delta\theta_{e\gamma}$	23 mrad

- The resolutions determine the **accidental background**
- For a given background we choose **R( $\mu$ )** and **running time**.
  - $\text{BG} = 0.5$  events
  - $\text{R}(\mu) = 1.2 \times 10^7 \mu/\text{sec}$
  - $T = 3.5 \times 10^7 \text{ sec}$  (2 years running time)
  - $\Rightarrow \text{SES} = 6 \times 10^{-14}$  ( $1.7 \times 10^{13}$  muons observed)
- NO candidate  $\Rightarrow \text{BR}(\mu \rightarrow e\gamma) < 1.2 \times 10^{-13}$  @ 90% CL
- Unlikely fluctuation (4 events)  $\Rightarrow \text{BR}(\mu \rightarrow e\gamma) \approx 2.4 \times 10^{-13}$

# Summary and Time Scale

- The experiment may provide a clean indication of **New Physics** or a strong constrain on SM extensions
- Measurements and detector simulation make us confident that we can reach the **SES of  $6 \times 10^{-14}$**  to  $\mu \rightarrow e\gamma$  ( $BR = 1.2 \cdot 10^{-13}$ )
- Final prototypes af (almost) all subdetectors were measured
  - Liquid Xe calorimeter Large Prototype
  - Timing counters
- Detector **assembly** during end 2006 **engineering run** end 2006 - beg. 2007



## Plans

- Data taking from 2007 on to reach  $10^{-13}$  sensitivity (90% CL)
- Obtain a “significant” result before the LHC era
- Eventual reach  $10^{-14}$  during LHC era